

## ***RESULTS OF A COCHRANE REVIEW***

### **ORAL GALACTOGOGUES (NATURAL THERAPIES OR DRUGS) FOR INCREASING BREAST-MILK PRODUCTION IN MOTHERS OF NON-HOSPITALIZED TERM INFANTS**

**Authors:** Siew Cheng Foong, May Loong Tan, Wai Cheng Foon, Lisa Marasco, Jacquelin Ho, Joo Howe Ong

**Study types** included: Randomized Controlled Trials (RCTs) and Quasi-RCTs

**Participants** Mothers breastfeeding or expressing for term healthy infants <6 mos of age

#### **Intervention Comparisons:**

1. Pharmacological oral galactagogue vs Placebo or no treatment
2. Natural oral galactagogue vs Placebo or no treatment
3. Galactagogue vs another Galactagogue

#### **Primary Outcome measures**

- Proportion of infants breastfeeding (excl or any) at 3, 4 and 6 mos postpartum
- Infant weight in trials where infants are receiving only own mothers' milk
- Volume of human milk measured in a specified amount of time


#### **Secondary Outcome measures**

- Adverse effects in mother or baby
- Ability of parent to reduce or stop formula supplementation
- Parental psychological status: satisfaction scores, depression scale, etc.

**SEARCH STRATEGY:** Standard search of Cochrane data base for Pregnancy and Childbirth group, derived from: Trial Registers, Medline, Embase, CINAHL, searches of 30 journals and proceedings of major conferences, weekly awareness alerts for 44 journals plus BioMed Central alerts., regional & content-specific databases; regional and content specific data bases; Also HERDIN (Philippines) and Napralert with special search terms. Secondary sources included following references, personal article collection of author LM. No language, geographic or date restrictions.

#### **Excluded studies:**

- **Pharmacological:** arginine aspartate (1); domperidone (6); growth hormone (3); iodine (3); luteotropin (1); metoclopramide (7); metoclopramide, domperidone and ferolactan (1); obron multivitamin (1); orgametril (1); oxytocin (9); oxytocin & sulpiride (1); Oestrogen and progestogen (1); Pitocin (1); pseudophedrine (1); sulpiride (1); thyrotropin releasing hormone (1).
- **Botanical:** Chasteberry (3); collagen soup (1); fenugreek (3); fenugreek/garlic mix (1); goat's rue (2); goat's rue/silymarin (1); garlic (1); glutamic acid (1); hedge nettle (3); humana still-tea (1); kyuki-choketu (2); Lactare (6); Leptaden (8); moringa (1); nutrition supplement (1); milk and eggs (1); Motherlove More Milk Plus (1); Mu-ying-li (1); Oligoplex/vitex (2); pectin extract (1); molocco placental extract (1); sesame (1); shatavari (1); Torbangun (2); various Japanese medicines (1); Yangxueshengru oral liquor (1)




### Characteristics of included studies

- **Locations:** North America (1); Latin America (3); Europe (3); Eurasia (6); East Asia (8); West Asia/Africa (4); South Asia (6); Southeast Asia (13)
- **Participants:** C-section only (2); Late term pregnancy only (1); Primips only (5); Female babies only (1); Infants 0-14d (21); Lactation deficient (18); older infants 2wks-6 mos (16); working mothers separated 8 hrs (1); (No milk problems (?);
- **Pharmacological interventions (4):** domperidone (2); metoclopramide (3); sulpiride (3); TRH (2);
- **Natural interventions (29):** Shatavari capsules (1); Cui Ru soup (1); Fennel tea (1); Fenugreek tea (1); Lactogenic food menu (1); Ginger capsules (1); Cottonseed capsules (1); Humana Still-Tee (2); Ixbut infusion (1); Malunggay/moringa capsules (4); Shatavari combo capsules (1); Pork leg soup (1); Shirafza drops (10); Silymarin sachet (1)
- **Galactagogue vs Galactagogue:** Domperidone vs malunggay (1); Torbangun vs fenugreek vs molocco (1); Mu Er Wu You vs Kun Yuna tong ru fu ye soup (1); Ru quan chong ji vs shengruzhi soup; fenugreek vs palm dates (1); fennel vs fenugreek black teas (1); Chanbao vs bu xue shen ru vs no intervention (1)


### Studies by outcome:

- Duration of breastfeeding: (3)
- Volume of milk measured (28)
  - By pre- and -post feeds (7)
  - By pre- and post feeds + residual milk (2)
  - Expression of milk by hand or pump (11)
  - Changes in breast dimensions (1)
  - Lai 4-hr pumping method (1)


- Infant weight (9) measured in various ways (total gain, weekly gain, mean % gain, etc)
- Adverse effects (17, but only 3 pre-specified)
- Ability to reduce supplementation (6)
- Maternal psychological status (4)




**Risk of bias graph:** review of authors’ judgements about each risk of bias item presented as percentages across all included studies



## Comparison 1.1 Infant Gain




## Comparison 1.2: Milk Volume




## Summary of Comparison 1

Pharmacological oral galactagogues compared to placebo or no treatment for increasing breast milk production in mothers of non-hospitalised term infants						
Patient or population: increasing breast milk production in mothers of non-hospitalised term infants						
Setting: Community						
Intervention: pharmacological oral galactagogues						
Comparison: placebo or no treatment						
Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	№ of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with placebo or no treatment	Risk with pharmacological oral galactagogues				
1: Proportion of infants bfg @ 3, 4, 6 mos		-	-	-	-	No studies reported this outcome
2: Infant weight (metoclopramide)	The mean infant weight was 0	MD 23 grams higher (47.71 lower to 93.71 higher)	-	20 (1 RCT)	LOW <sup>1</sup>	
3: Milk Volume (all galactagogues)	The mean milk volume was 0	MD 63.82 mL higher (25.91 higher to 101.72 higher)	-	151 (3 RCTs)	LOW <sup>1</sup>	
4: Adverse Effects (mother or infant)	With metoclopramide, tiredness, headache, nausea were reported more frequently than placebo group. With domperidone dry mouth were reported more frequently than placebo group and no extrapyramidal effects reported. With sulpiride tiredness and headache were reported more frequently than placebo group. No adverse effects reported with thyrotropin-releasing hormone.		-	133 (5 RCTs)	VERY LOW <sup>2,3</sup>	Adverse effects were generally poorly reported.
*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
CI: Confidence interval; RR: Risk ratio; OR: Odds ratio						
GRADE Working Group grades of evidence						
High certainty: We are very confident that the true effect lies close to that of the estimate of the effect						
Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different						
Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect						
Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect						


## Comparison 2.1: Infant weight



## Comparison 2.3: Volume of supplement




## Comparison 2.2: Milk volume





## Comparison 2.4-2.6



## Comparison 2 Summary

2 Natural oral galactagogues compared to placebo or no treatment for increasing breast milk production in mothers of non-hospitalised term infants

Natural oral galactagogues compared to placebo or no treatment for increasing breast milk production in mothers of non-hospitalised term infants						
Patient or population: increasing breast milk production in mothers of non-hospitalised term infants						
Setting: Community						
Intervention: natural oral galactagogues						
Comparison: placebo or no treatment						
Outcomes	Anticipated absolute effects <sup>*</sup> (95% CI)		Relative effect (95% CI)	N <sup>o</sup> of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with placebo or no treatment	Risk with natural oral galactagogues				
1: % of infants bfg @ 3, 4, 6 mos	One study reported no significant difference in breastfeeding rates at six months for Mother's Milk Tea.		-	72 (1 RCT)	VERY LOW <sup>1,2</sup>	
2: Infant weight (metoclopramide)	see comment	see comment	-	275 (3 RCTs)	VERY LOW <sup>3,4</sup>	Only subgroup analyses per galactagogue type. No meta-analysis.
3: Milk Volume (all galactagogues)	see comment	see comment	-	962 (13 RCTs)	VERY LOW <sup>3,4,5</sup>	Only subgroup analyses per galactagogue type. No meta-analysis.
4: Adverse Effects (mother or infant)	Almost all reported "no adverse effects." Where reported, they were single occurrences such as maple-syrup urine with fenugreek, ginger and turmeric mix; nausea and urticaria in infants with Shiratza.		-	(10 RCTs)	VERY LOW <sup>2,7</sup>	Adverse effects were poorly reported.
*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
CI: Confidence interval; RR: Risk ratio; OR: Odds ratio						
GRADE Working Group grades of evidence						
High certainty: We are very confident that the true effect lies close to that of the estimate of the effect						
Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different						
Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect						
Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect						

## HETEROGENEITY OF STUDIES:

### Study participants:

- ❖ Different ages of babies enrolled
- ❖ Breastfeeding routines frequently not described
- ❖ Normal supply vs low supply; causes of low supply
- ❖ Parity, c-section, etc.

### Interventions and methodologies

- ❖ Dosages
- ❖ Treatment durations
- ❖ Natural botanical materials rarely validated
- ❖ Questionable placebos

### Outcome measurements

- ✓ Using baby's gain- sometimes rate, sometimes weight
- ✓ Using milk transfer as a proxy for production- what about residual?
- ✓ Time points of measurement sometimes different for the groups

### **Insights:** *Rescue Vs Proactive Study Goals*

## **CONCLUSION**

- ✓ *Very low certainty evidence that oral galactagogues in the review might improve infant weight and milk volume*
- ✓ *We are uncertain if one galactagogue is better than another, or their use would result in any harm*
- *Due to substantial heterogeneity of the studies, imprecision of measurements and incomplete reporting, we are very uncertain about the magnitude of the effect*
- *Very little available evidence of effect on bfg rates at 3, 4 or 6 mos*
- *Not enough evidence to judge harm*
- *Management should always be the first step before employing galactagogues=*

### **For quality evidence, we need:**

- *High quality RC*
- *Set of core outcomes to standardize measurements*
- *Strong basis for dosages and forms used*



## **HOW TO BUILD A BETTER STUDY**

- ✓ Methodology and reporting need to be at same academic standard as other pharmaceutical interventions
- ✓ Mandatory lactation support provided
- ✓ Infants of similar ages
- ✓ For botanicals, validation of material and purity
- ✓ Preparation of plant material described
- ✓ Rationale of form and dosage
- ✓ Measurement of milk by volume should include transfer & residual expressed, preferably 24 hrs
- ✓ Report total # of breastfeeding/expression sessions
- ✓ Report duration of 'any' and 'exclusive' bfg to 6 mos, as that is the ultimate goal
- ✓ Identify, screen for and report side and adverse effects in mother and baby

### **Priority for future Studies:**

- For women with IMS, attempt to identify etiology of low production
- Test more commonly used galactagogues first
- Test multiple dosages to determine most effective therapeutic dosage

### **Priority – Related research needs**

- Determine a standard for defining lactation insufficiency beyond maternal perception
- Determine a standard method to measure “milk volume”, including measure tools and duration of measurement. Explore usefulness of Lai method
- Determine mechanisms by which a galactagogue may increase milk production. Take clues from animal studies. This may lead to better rationales for choosing one galactagogue over another

## Bibliography

### Included studies:

- Aono T, Aki T, Koike K, Kurachi K. Effect of sulpiride on poor puerperal lactation. *American Journal of Obstetrics and Gynecology* 1982;143:927-32
- Balahibo M, Balde J, Baldovino Z, Baquiran P, Barillos C, Barredo C, Madarcos F, Mendoza S. A randomized, doubleblinded parallel-controlled clinical trial on the effectiveness of different doses of *Moringa oleifera* (Malunggay) in promoting growth in infants of breastfeeding mothers from UERMMMC and different hospitals in Metro Manila from June 2000 to January 2001. *UERMMMC Journal of Health Sciences* 2001;5(1):21-26.
- Barguno JM, Del Pozo E, Cruz M, Figueras J. Failure of maintained hyperprolactinemia to improve lactational performance in late puerperium. *Journal of Clinical Endocrinology and Metabolism* 1988;66:876-9
- Briton Medrano G, Perez L. The efficacy of malunggay (*moringa oleifera*) given to near term pregnant women in inducing early postpartum breast milk production- a double blind randomized clinical trial. 2002.
- Chan ML. Effect of Ixbut consumption euphorbia lancifolia schlecht about density and volume of breastmilk [Efecto del consumo de Ixbut Euphorbia lancifolia schlecht sobre la densidad y el volumen de la leche materna]
- Damanik R, Wahlqvist ML, Wattanapenpaiboon N. Lactagogue effects of Torbangun, a Batakese traditional cuisine. *Asia Pacific Journal of Clinical Nutrition* 2006;15(2):267-74.
- De Gezelle H, Ooghe W, Thiery M, Dhont M. Metoclopramide and breast milk. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 1983;15:31-6.
- Di Pierro F, Callegari A, Carotenuto D, Tapia MM. Clinical efficacy, safety and tolerability of BIO-C (micronized Silymarin) as a galactagogue. *Acta Bio-Medica de l Ateneo Parmense* 2008;79(3):205-10.
- Espinosa-Kuo CL. A randomized controlled trial on the use of malunggay (*Moringa oleifera*) for augmentation of the volume of breastmilk among mothers of term infants. *The Filipino Family Physician* 2005;43(1):26-33.
- Fang Q, He XH, Zhang ZF. Clinical observation on treatment of postnatal lack of lactation by Ruquan Drink. *China Journal of Chinese Materia Medica [Zhong Guo Zhong Yao Za Zhi]* 2003;28(8):792-3
- Ghasemi V, Kheirkhah M, Samani LN, Vahedi M. The effect of herbal tea containing fennel seed on breast milk sufficiency signs and growth parameters of Iranian infants. *Shiraz E Medical Journal* 2014;15(4):1-5.
- Gupta M, Shaw B. A double-blind randomized clinical trial for evaluation of galactagogue activity of asparagus racemosus willd. *Iranian Journal of Pharmaceutical Research* 2011;10(1):167-72.
- Huang HL, Li SL, Li LZ. Clinical study of Xian Tong Ru oral liquor in enhancing early lactation. <http://www.cnki.net> 2000; 9(23):2330-2331.
- Inam I, Hashmi AB, Shahid A. A comparison of efficacy of domperidone and placebo among postnatal women with inadequate breast milk production. *Pakistan Journal of Medical and Health Sciences* 2013;7(2):1-3.
- Jantarsaengaram S, Sreewapa P. Effects of domperidone on augmentation of lactation following cesarean delivery at full term. *International Journal of Gynecology and Obstetrics* 2012;116(3):240-3.
- Jiang J, Jiang H, Jiang R. Clinical Study of Chanbao Oral Liquid on Stimulating Milk Secretion. *World Journal of Integrated Traditional and Western Medicine* 2006;1(2):114-6.
- Kaupila A, Anunti P, Kivinen S, Koivisto M, Ruokonen A. Metoclopramide and breast feeding: efficacy and anterior pituitary responses of the mother and the child. *European Journal of Obstetrics & Gynecol and Reproductive Biology* 1985; 19:19-22.
- Li DM, Liu YQ, Li LM. Observation of Muer Wuyou decoction on postpartum hypogalactia. *Hebei Journal of Traditional Chinese Medicine* 2010;32(5):655-7.
- Manjula S, Sultana A, Rahman K. Clinical efficacy of *Gossypium herbaceum* L. seeds in perceived insufficient milk (PIM) supply: A randomized single-blind placebo-controlled study. *Oriental Pharmacy and Experimental Medic* 2014; 14(1):77-85.
- Paritakul P, Ruangrongmorakot K, Laosooksathit W, Suksamarnwong M, Puapornpong P. The Effect of Ginger on Breast Milk Volume in the Early Postpartum Period: A Randomized, Double-Blind Controlled Trial. *Breastfeeding medicine : the official journal of the Academy of Breastfeeding Medicine* 2016;11(7):361-5. [ PubMed: 27505611]
- Sakha K, Behbahan AG. Training for perfect breastfeeding or metoclopramide: which one can promote lactation in nursing mothers? *Breastfeeding Medicine* 2008;3(2):120-3.
- Abeer El Sakka, Mostafa Salama, Kareem Salama. The Effect of Fenugreek Herbal Tea and Palm Dates on Breast Milk Production and Infant Weight. *Journal of Pediatric Sciences* 2014;6:e202. [DOI: <http://dx.doi.org/10.17334/jps.30658>]

- Shariati M, Mammoori GH, Khadivarzadeh T. The Effect of Taking "Shirafza Drops" by Mothers on the Weight Gain of Infants up to the Age of Six Months. *Journal of Sabzevar School of Medical Sciences* 2004;11(3(33)):20-26.
- Sharma S, Ramji S, Kumari S, Bapna JS. Randomized controlled trial of Asparagus racemosus (Shatavari) as a lactagogue in lactational inadequacy. *Indian Pediatrics* 1996;33:675-7.
- Su WQ, Chen YJ. Clinical Observation on Effect of Cuiru Decoction in the Treatment of Postpartum Hypogalactia. *Chinese Journal of Medicinal Guide* 2008;10(3):420-1.
- Sy W, Bernardo C. A comparative study of Malunggay (*Moringa oleifera*) and Domperidone as a galactagogue among mothers of term infants. *Herdin Record #*: NCR-TMC-14021916374880 20
- Thaweekul P, Thaweekul Y, Sritipsukho P. The efficacy of hospital-based food program as galactagogues in early period of lactation. *Journal of the Medical Association of Thailand* 2014;97(5):478-82.
- Tirak U, Dilli D, Emeksiz S, Dallar Y. The Effect of a Galactagogue Herbal Tea on Weight Gain During Newborn Period [Galaktagog Bir Bitki Çayın Yenidoğan Döneminde Tart Alm Üzerine Etkisi]. *Çocuk Dergis* 2008;8(1):14-20.
- Turkylmaz C, Onal E, Hirfanoglu IM, Turan O, Koc E, Ergenekon E, et al. The effect of galactagogue herbal tea on breast milk production and short-term catch-up of birth weight in the first week of life. *Journal of Alternative and Complementary Medicine* 2011;17(2):139-42.
- Wagner CL, Morella K, Boan AD, Guille C, Marzolf A, Finch CW, et al. The safety of a Mother's Milk Tea (MMT): Results of a randomized controlled trial. In: *Ped Acad Societies Annual Meeting*; 2016 April 30-May 3; Baltimore, USA. 2016:2160.8.
- Xu JH, Gao QJ. Effect of Galactagogue on the Lactation and Infant Body Weight in Delivery Women [J]. *Journal of Nursing Science* 2000;11:646-7.
- Yabes-Almirante C, Lim CHTN. Effectiveness of Natalac as a galactagogue. *Journal of the Philippine Medical Association* 1996;71(3):265-72.
- Ylikorkala O, Kaupila A, Kivinen S, Viinikka L. Sulpiride improves inadequate lactation. *BMJ* 1982;285:249-51.
- Yong Y, Shanling T, Chengye Y. Effect of ShengRu Mistura on the Serum and Milk Contents of PRL in Postpartum Women. *Journal of Radioimmunology* 2005;18(6):467-8.
- Zarate A, Villalobos H, Canales E, Soria J, Arcovedo F, MacGregor C. The effect of oral administration of thyrotropin-releasing hormone on lactation. *Journal of Clinical Endocrinology and Metabolism* 1976;43:301-5.

### **Other references:**

- Edwards, R. (2018). An Exploration of Maternal Satisfaction With Breastfeeding as a Clinically Relevant Measure of Breastfeeding Success. *J Hum Lact*, 34(1), 93-96. doi:10.1177/0890334417722509
- Flaherman, V. J., Hicks, K. G., Cabana, M. D., & Lee, K. A. (2012). Maternal experience of interactions with providers among mothers with milk supply concern. *Clin Pediatr (Phila)*, 51(8), 778-784. doi:10.1177/0009922812448954
- Gagnier, J., Boon, H., Rochon, P., Barnes, J., Moher, D., Bombardier, C., & Group, C. (2006). Improving the quality of reporting of randomized controlled trials evaluating herbal interventions: implementing the CONSORT statement [corrected]. *Explore (NY)*, 2(2), 143-149. doi:10.1016/j.explore.2005.12.003
- Gagnier, J., & Moher, D. (2006). Helping consumers of healthcare knowledge: Improving the quality of reports of randomized controlled trials. *Explore (NY)*, 2(2), 141-142. doi:10.1016/j.explore.2005.12.010
- Gagnier, J. J., Boon, H., Rochon, P., Moher, D., Barnes, J., Bombardier, C., & Group, C. (2006). Recommendations for reporting randomized controlled trials of herbal interventions: Explanation and elaboration. *J Clin Epidemiol*, 59(11), 1134-1149. doi:10.1016/j.jclinepi.2005.12.020
- Gagnier, J. J., Boon, H., Rochon, P., Moher, D., Barnes, J., Bombardier, C., & Group, C. (2006). Reporting randomized, controlled trials of herbal interventions: an elaborated CONSORT statement. *Ann Intern Med*, 144(5), 364-367.
- Gagnier, J. J., DeMelo, J., Boon, H., Rochon, P., & Bombardier, C. (2006). Quality of reporting of randomized controlled trials of herbal medicine interventions. *Am J Med*, 119(9), 800 e801-811. doi:10.1016/j.amjmed.2006.02.006
- Labbok, M. H., & Starling, A. (2012). Definitions of breastfeeding: call for the development and use of consistent definitions in research and peer-reviewed literature. *Breastfeeding Medicine*, 7(6), 397-402.
- Lauer, J. A., Betran, A. P., Victora, C. G., de Onis, M., & Barros, A. J. (2004). Breastfeeding patterns and exposure to suboptimal breastfeeding among children in developing countries: review and analysis of nationally representative surveys. *BMC Med*, 2, 26. doi:10.1186/1741-7015-2-26
- Noel-Weiss, J., Boersma, S., & Kujawa-Myles, S. (2012). Questioning current definitions for breastfeeding research. *International breastfeeding journal*, 7(9).
- Organization, W. H., & Unicef. (2003). Global strategy for infant and young child feeding.
- Spatz, D., & Lessen, R. (2011). Risks of not breastfeeding. *Morrisville, NC: International Lactation Consultant Association*.